

AMENABILITY OF THE ALGEBRAS $R(S)$, $F(S)$ OF A TOPOLOGICAL SEMIGROUP S

MASSOUD AMINI, ALIREZA MEDGHALCHI

ABSTRACT. For a locally compact Hausdorff semigroup S , the L^∞ representation algebra $R(S)$ was extensively studied by Dunkl and Ramirez. The Fourier-Stieltjes algebra $F(S)$ of a topological semigroup was studied by Lau. The aim of this paper is to investigate these two algebras and study the amenability of them with respect to the structure of S .

1. INTRODUCTION.

Dunkl and Ramirez defined the subalgebra $R(S)$ of the algebra of weakly almost periodic functions on S , $WAP(S)$ [2]. This algebra is called the L^∞ -representation algebra of S . In fact $R(S)$ is the set of all functions $f(x) = \int_X (Tx)gd\mu$ ($x \in S$) where (T, X, μ) is an L^∞ -representation of S and $g \in L^1(X, \mu)$.

In [3] Lau studied the subalgebra $F(S)$ of $WAP(S)$ of a topological semigroup S with involution. If S is commutative, then $F(S) \subseteq R(S)$ and in particular, if G is an abelian topological group, then $F(G) = R(G) = \widehat{M(\hat{G})}$ where $\hat{G}$ is the dual group of G . If S is a topological* - semigroup with an identity, then $F(S)$ is the linear span of positive definite functions on S . By ([3], theorem 3.2) $F(S)$ is a subalgebra of $WAP(S)$. In this paper, we investigate the structure of $R(S), F(S)$

1991 *Mathematics Subject Classification.* 43A37, 43A20.

Key words and phrases. L^∞ -representation algebra, Fourier Stieltjes algebra, amenability, weakly and strongly almost periodic functions.

This research was partly supported by IPM.

and $\overline{R(S)}, \overline{F(S)}$, the normed closures of $R(S)$ and $F(S)$ and show that these are left introverted subalgebras of $\text{WAP}(S)$ and study left (resp. right) amenability of $R(S)$, $F(S)$, $\overline{R(S)}$, $\overline{F(S)}$. Also we show that $\text{SAP}(S) \subseteq \overline{F(S)}$, where $\text{SAP}(S)$ is the Banach algebra of strongly almost periodic functions on S , and then show that $\overline{F(S)}$ is left amenable if and only if $\overline{F(S)} = \text{SAP}(S) \oplus C$, where C is a closed linear subspace of $\overline{F(S)}$.

2. PRELIMINARIES

Let S be a topological semigroup. Let λ be a probability measure on a measurable space X . We know that $L^\infty(X, \lambda)$ is a commutative W^* -algebra. An L^∞ -representation (T, X, μ) of S is a weak*-continuous homomorphism T of S into the unit ball of $L^\infty(X, \mu)$.

Definition 2.1. The representation algebra $R(S)$ is the set of all functions f such that $f(x) = \int_X T_x(g) d\mu$, where (T, X, μ) is an L^∞ -representation of S and $g \in L^1(X, \mu)$.

Proposition 2.2. $R(S)$ is a normed subalgebra of $\text{WAP}(S)$ with pointwise multiplication. It is closed under conjugation, translation invariant and contains constant functions. It is complete with its norm.

Proof. See Theorem 1.6 and Proposition 1.4 of [2].

Following ([1] pages 72, 77 and section 3) we have the following definitions:

Definition 2.3. Let F be a translation invariant subalgebra of $B(S)$ (the space of all bounded functions on S). F is said m -left introverted if $T_\mu F \subseteq F$ ($\mu \in MM(F)$) where $T_\mu f(x) = \mu(l_x f)$, $l_x f(y) = f(xy)$ and $MM(F)$ the set of all multiplicative means on F .

Definition 2.4. An admissible subalgebra of $B(S)$ is a normed closed, conjugate closed, translation invariant, left m -intrverted subalgebra of $B(S)$ containing constant functions.

Following Lau we have the following definition [3].

Definition 2.5. Let S be a topological semigroup with continuous involution $*$. Let M be a W^* -algebra and $M_1 = \{x \in M \mid \|x\| \leq 1\}$. By a $*$ -representation of S we mean (ω, M) , where ω is a $*$ -homomorphism of S into M_1 . Let $\Omega(S)$ be the set of all σ -continuous $*$ -representations of S such that $\overline{\omega(S)}^\sigma = M$. Let $F(S)$ be the set of all functions f on S such that $f = \hat{\psi}$ for some $\psi \in M_*$ (predual of M), where $\hat{\psi} = \psi \circ \omega$. Let $f \in F(S)$, in [3] Lau defines $\|f\|_\Omega = \inf\{\|\psi\| \mid \psi \in M_*, \hat{\psi} = f \text{ for some } (\omega, M) \in \Omega(S)\}$. Note that $\sigma = \sigma(M, M_*)$.

Following [3] we have the following proposition.

Proposition 2.6. $F(S)$ is a commutative subalgebra of $WAP(S)$ which is conjugate closed, translation invariant, containing the constant functions. Also, if $f \in F(S)$, then $f^* \in F(S)$, where $f^*(s) = \overline{f(s^*)}$. Furthermore, $\|\cdot\|_\Omega$ is a norm on $F(S)$ and $(F(S), \|\cdot\|_\Omega)$ is a commutative normed algebra with unit.

Proof. This is theorem 3.2 of [3] where its proof was referred to [2].

3. AMENABILITY OF $F(S), R(S), \overline{F(S)}, \overline{R(S)}$.

In this section we assume that S is a topological semigroup with a continuous involution.

In the following theorem $K(X), K(Y)$ are the minimal ideals of X, Y , respectively.

Theorem 3.1. Let $Y = \sigma(F(S))$ (the spectrum of $F(S)$) which is a semigroup by proposition 5.4 in [3]. Then $F(S)$ is amenable if and only if $K(Y)$ is a topological group.

Proof. Assume $F(S)$ is amenable and let $\overline{F(S)}$ be the norm closure of $F(S)$ in $\text{WAP}(S)$. Then clearly $\overline{F(S)} \subseteq \text{WAP}(S)$. By (theorem 3.2 [3]), $\|f\|_\infty \leq \|f\|_\Omega$ ($f \in F(S)$). So the embedding function $i : F(S) \longrightarrow \overline{F(S)}$ is continuous. Therefore by (theorem 1.9. [5]), $\overline{F(S)}$ is amenable. Now again by (theorem 3.2, [1]) $F(S)$ is a translation invariant, conjugate closed subalgebra of $\text{WAP}(S)$ and contains constant functions, hence $\overline{F(S)}$ is norm closed, translation invariant, conjugate closed subalgebra of $\text{WAP}(S)$ containing constant functions. Therefore by (corollary 4.2.7, [1]) $\overline{F(S)}$ is introverted. So, it is an m -admissible subalgebra of $\text{WAP}(S)$. Hence, by (theorem 4.2.14, [1]) $K(X)$ is a topological group, where X is $S^{\overline{F(S)}}$, the maximal ideal space of $\overline{F(S)}$. Now let $\pi = i^* : \overline{F(S)}^* \longrightarrow F(S)^*$ is the adjoint mapping of i . Clearly π is norm decreasing and onto and $\pi(X) = Y$. Thus $\pi(K(X)) = K(Y)$ is a topological group. Conversely, if $K(Y)$ is a topological group, then $K(X)$ is a topological group, so by (4.2.14, [1]) $\overline{F(S)}$ is amenable, and so is $F(S)$.

We have another consequence of (theorem 4.2.14, [1]). In fact, $\overline{F(S)}$ is left (right) amenable if and only if $K(X)$ is a minimal right (left) ideal of X .

Now, we extend the above theorem for left (resp. right) amenability of $F(S)$.

Theorem 3.2. $F(S)$ is left (right) amenable if and only if $K(Y)$ is minimal right (left) ideal of Y .

Proof. Let $F(S)$ be left amenable and $f \in \overline{F(S)}$. Then there is a sequence $\{g_n\}$ in $F(S)$ such that $g_n \xrightarrow{\|\cdot\|_\infty} f$. Clearly $\{m(g_n)\}$ is a Cauchy sequence in $\overline{F(S)}$, so it is convergent. We define $\bar{m}(f) = \lim_{n \rightarrow \infty} m(g_n)$. Since $\bar{m}$ is positive, $\bar{m}(1) = 1$

and $\bar{m}(l_x f) = \lim_{n \rightarrow \infty} m({}_x g_n) = \lim_{n \rightarrow \infty} m(g_n) = \bar{m}(f)$, it follows that $\bar{m}$ is a left invariant mean on $\overline{F(S)}$. Hence, $\overline{F(S)}$ is left amenable. By (theorem 4.2.14, [1]) $K(X)$ is a minimal right ideal of X . Now, we have $\pi(K(X)) = K(Y)$, therefore by (corollary 1.3.16, [1]) $K(Y)$ is a minimal right ideal of Y . Conversely, if $K(Y)$ is a minimal right ideal of Y , then by the same argument $K(X)$ is a minimal right ideal of X and therefore, again by (theorem 4.2.14, [1]) $\overline{F(S)}$ is left amenable. Thus $F(S)$ is left amenable. The right version is proved in a similar way.

Next we show that $F(S)$ is an F -algebra in the sense of Lau [4].

Proposition 3.3. If S is a unital topological semigroup with continuous involution, then $F(S)$ is an F -algebra.

Proof. Each element of $F(S)$ is of the form $\hat{\psi} = \psi \circ \omega_\Omega$, for some $\psi \in (M_\Omega)_*$, where $(\omega_\Omega, M_\Omega)$, is the universal representation of S [3]. Take $\psi_1, \psi_2 \in (M_\Omega)_*$, then by the Gelfand-Naimark-Segal construction, there are (unit) vectors $\xi_i, \eta_i \in H_\Omega$ such that $\psi_i(x) = \langle x \xi_i, \eta_i \rangle$ ($x \in M_\Omega, i = 1, 2$). Then

$$\hat{\psi}_1 \hat{\psi}_2(s) = \hat{\psi}_1(S) \hat{\psi}_2(s) = \langle \omega_\Omega(s) \xi_1, \eta_1 \rangle \langle \omega_\Omega(s) \xi_2, \eta_2 \rangle = \langle \omega_\Omega(s) \xi_1 \otimes \xi_2, \eta_1 \otimes \eta_2 \rangle \quad (s \in S).$$

Hence, if $1 \in W_\Omega^*(S)$ is the identity element (see [3] for the notation), then

$$\langle 1, \hat{\psi}_1 \hat{\psi}_2 \rangle = \langle \xi_1 \otimes \xi_2, \eta_1 \otimes \eta_2 \rangle = \langle \xi_1, \eta_1 \rangle \langle \xi_2, \eta_2 \rangle = \langle 1, \hat{\psi}_1 \rangle \langle 1, \hat{\psi}_2 \rangle,$$

and we are done.

Corollary 3.4. If S is a unital topological semigroup with continuous involution, then $W_\Omega^*(S)$ has a topological left invariant mean.

Proof. This follows from above proposition and (theorem 4.1, [4]). Note that $F(S)$ is always left amenable in the sense of [4].

Remark 3.5. (a) If S is an idempotent commutative topological semigroup with involution $s = s^*$, then by (3.3(c), [3]) $F(S) = R(S)$. Therefore in this special case the above results hold for $R(S)$ too.

(b) By Proposition 2.2, $R(S)$ is right translation invariant, conjugate closed, containing constant functions. Therefore when S is commutative, $\overline{R(S)} \subseteq \text{WAP}(S)$ is norm closed, translation invariant, conjugate closed subalgebra of $\text{WAP}(S)$ containing constant functions. therefore $\overline{R(S)}$ is an m -admissible subalgebra of $\text{WAP}(S)$ (theorem 4.2.14, [1]). Hence, the results of theorems 3.1 and 3.2 hold for $\overline{R(S)}$.

(c) If G is a non-compact locally compact abelian group, then $\text{WAP}(G) \neq \overline{R(G)} = \overline{M(\hat{G})}$ (see 52.10, [2]), or even if S is a non compact locally compact subsemigroup of a locally compact group, then $\text{WAP}(S) \neq \overline{R(S)}$ (see (5.2.12, [2])).

(d) Example 4.2 of [3] also shows that $\overline{F(S)} \neq \overline{R(S)}$ may happen.

(e) When S is an abelian semigroup with involution, we have another interesting result. By (4.3.8, [1]) $\text{SAP}(S)$ (the Banach space of all strongly almost periodic functions on S) is the closed linear span of characters. Therefore, by (3.3.(e), [1]) $\text{SAP}(S) \subseteq F(S) \subseteq \text{WAP}(S)$.

Now by (theorem 4.3.12, [1]) $\overline{F(S)}$ is left amenable if and only if $\overline{F(S)} = \text{SAP}(S) \oplus C$ where C is a left translation invariant, closed linear subspace of $\overline{F(S)}$, if and only if there exists a projection of $\overline{F(S)}$ onto $\text{SAP}(S)$ that commutes with left translations. Since, the amenability of $F(S)$ and $\overline{F(S)}$ are equivalent, the amenability of $F(S)$ is equivalent to each of the above conditions.

(f) If S is a topological group, then $Y = \sigma(B(S))$ is a σ -weakly compact, self adjoint subset of the unit ball of $W^*(S)$ and $Y \cap W^*(S)_u \simeq S$ (see [6] for details and notations). If X is $\overline{B(S)}$ -compactification of S and $G_1 = K(Y)$, $G_2 = K(X)$, then by theorem 3.1. and (3.4, [3]), the Banach algebras $A(S)$ and $B(S)/A(S)$ are

both amenable iff any of G_1 or G_2 is a topological group. This is interesting in the light of the fact that we don't know exactly when $A(S)$ is amenable.

Next we show that remark (e) indeed applies to any topological $*$ -semigroup.

Lemma 3.6. Let S be a topological $*$ -semigroup, then

$$\text{SAP}(S) \subseteq \overline{F(S)} \subseteq \text{WAP}(S).$$

Proof. Let $\{\pi, H\}$ be a (finite dimensional) unitary representation of S , $\xi, \eta \in H$, and $u(s) = \langle \pi(s)\xi, \eta \rangle$ ($s \in S$). Put $M = \langle \pi(S) \rangle^{-\sigma} \subseteq B(H)$ and $\alpha = (\pi, M) \in \Omega(S)$. Consider $\psi(x) = \langle x\xi, \eta \rangle$ ($x \in M$). Then obviously $\psi \in M_*$. Indeed ψ is w^* -continuous on $B(H)$: if $\{T_\alpha\} \subseteq B(H)$ and $T_\alpha \xrightarrow{w^*} T$ in $B(H)$, then considering the rank one operator $\xi \otimes \eta$ in $B(H)_*$, we have

$$\langle T_\alpha \xi, \eta \rangle = \langle T_\alpha, \xi \otimes \eta \rangle \xrightarrow{\alpha} \langle T, \xi \otimes \eta \rangle = \langle T\xi, \eta \rangle$$

i.e. $\psi(T_\alpha) \longrightarrow \psi(T)$. Now we have $\hat{\psi} = \psi \circ \pi = u$, hence $u \in F(S)$. Therefore $\text{SAP}(S) \subseteq \overline{F(S)}$. The other inclusion is contained in [3].

Now the proof of the following theorem is exactly as in remark (e).

Theorem 3.7. Let S be a topological semigroup with continuous involution. Then the following are equivalent:

- a) $R(S)$ is amenable.
- b) $F(S)$ is amenable.
- c) $\overline{R(S)}$ is amenable.
- d) $\overline{F(S)}$ is amenable.
- e) $\overline{F(S)} = \text{SAP}(S) \oplus C$, where C is a translation invariant, closed linear subspace of $\overline{F(S)}$.

- f) There exists a projection of $F(S)$ onto $SAP(S)$ that commutes with translations.

REFERENCES

- [1] Berglund, John F., Junghenn, Hugo, D., Milnes, P., Analysis on semigroups, John Wiley & Sons, 1989.
- [2] Dunkl, Charles, F., Ramirez, Donald, E., Representation of commutative semitopological semigroup, Lecture Notes in Mathematics, 435(1975) Springer-Verlag.
- [3] Lau, A. T., The Fourier Stieltjes algebra of a topological semigroup with involution, Pacific J. Math. vol. 77, No. 1 (1978), 165-181.
- [4] Lau, A. T., Analysis on a class of Banach algebras with applications to harmonic analysis on locally compact groups and semigroups, Fundamenta Math. vol. 68 (1983), 161-175.
- [5] Pier, Jean. Paul, Amenable Banach algebras, John Wiley, 1988.
- [6] Walter, Martin E., W^* -algebras and nonabelian Harmonic analysis, J. Func. Anal. vol. 11 (1972), 17-38.

DEPARTMENT OF MATHEMATICS, SHAHID BEHESHTI UNIVERSITY, EVIN, TEHRAN 19839, IRAN,
M-AMINI@CC.SBU.AC.IR

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, 106 WIGGINS
ROAD, SASKATOON, SASKATCHEWAN, CANADA S7N 5E6, MAMINI@MATH.USASK.CA

DEPARTMENT OF MATHEMATICS, TEACHER TRAINING UNIVERSITY, TEHRAN, IRAN